

Computational Intelligence

Lecture 11: Linguistic Variables and Fuzzy Rules

Farzaneh Abdollahi

Department of Electrical Engineering


Amirkabir University of Technology

Linguistic Variables
Linguistic Hedge

Fuzzy IF-Then Rules
Fuzzy Proposition
Interpretation of Fuzzy Rules

Linguistic Variables

- ▶ **Linguistic Variable:** when a variable can take words in natural languages as its values.
 - ▶ the words are characterized by fuzzy sets defined in the universe of discourse
- ▶ **Example:** The weather temperature: a variable $x \in [-20 \ 40]$
 - ▶ Let's define three fuzzy sets "Cold", "Pleasant," and "Hot"
 - ▶ x : a linguistic variable, \rightsquigarrow "x is cold,"
 - ▶ x also can take numbers in $[-20 \ 40] \rightsquigarrow, x = 10^\circ \text{C}$



- ▶ **Linguistic Variable:** is characterized by (X, T, U, M)
 - ▶ **X:** the name of the linguistic variable (the weather temperature)
 - ▶ **T:** the set of linguistic values that X can take
 $(T = \{cold, pleasant, hot\})$
 - ▶ **U:** the actual physical domain in which the linguistic variable X takes its quantitative (crisp) values $(U = [-20, 40])$.
 - ▶ **M:** a semantic rule that relates each linguistic value in T with a fuzzy set in U ; (M relates "cold," "pleasant," and "hot" with the membership functions shown in Fig)

- ▶ **Linguistic Hedge:**
 - ▶ One may use more than one word to describe the ling. var. ("very cold", "not pleasant", "more or less hot")
 - ▶ The value of a ling. var. is a composite of atomic term $x = x_1 x_2 \dots x_n$
 - ▶ The atomic terms can be classified into:
 - ▶ **Primary terms:** labels of fuzzy sets; "cold," "pleasant," and "hot."
 - ▶ **Complement:** "not", "and" and "or."
 - ▶ **Hedges:** "very," "slightly," "more or less,"

Some Examples of Linguistic Hedge

- ▶ Let A be a fuzzy set in U
 - ▶ **very A** : a fuzzy set in U $\mu_{\text{very } A}(x) = [\mu_A(x)]^2$
 - ▶ **more or less A** : a fuzzy set in U $\mu_{\text{more or less } A}(x) = [\mu_A(x)]^{1/2}$
- ▶ **Example:** $U = \{10, \dots, 130\}$
 - ▶ fuzzy set "fast" = $1/130 + 1/120 + 0.8/100 + 0.6/80 + 0.1/40$
 - ▶ "very very fast" = $1/130 + 1/120 + 0.4096/100 + 0.1296/80 + 0.0001/40$
 - ▶ "more or less fast" = $1/130 + 1/120 + 0.8944/100 + 0.7746/80 + 0.3162/40$

Fuzzy IF-Then Rules

- ▶ **IF** <fuzzy proposition>, **THEN** <fuzzy proposition>
- ▶ There are two types of fuzzy proposition
 - ▶ **Atomic fuzzy proposition**: a single statement
 x is A
 - ▶ **Compound fuzzy proposition**: a composition of atomic fuzzy propositions using the connectives "and," "or," and "not"
 x is not S and x is not F
 - ▶ S, F, A are fuzzy sets
- ▶ Compound fuzzy propositions should be considered as fuzzy relations.

- ▶ Let x, y : linguistic variables in the physical domains U, V , and A, B : fuzzy sets in U, V
 - ▶ Connective "and" x is A and y is B use fuzzy intersection:
 - ▶ $A \cap B \in U \times V: \mu_{A \cap B}(x, y) = t[\mu_A(x), \mu_B(y)]$
 - ▶ $t: [0, 1] \times [0, 1] \rightarrow [0, 1]$ is any t-norm.
 - ▶ Connective "or" x is A or y is B use fuzzy union:
 - ▶ $A \cup B \in U \times V: \mu_{A \cup B}(x, y) = s[\mu_A(x), \mu_B(y)]$
 - ▶ $s: [0, 1] \times [0, 1] \rightarrow [0, 1]$ is any s-norm.
 - ▶ Connective "not" x is not A use fuzzy complements.

- ▶ Let x, y : linguistic variables in the physical domains U, V , and A, B : fuzzy sets in U, V
 - ▶ Connective "and" x is A and y is B use fuzzy intersection:
 - ▶ $A \cap B \in U \times V: \mu_{A \cap B}(x, y) = t[\mu_A(x), \mu_B(y)]$
 - ▶ $t: [0, 1] \times [0, 1] \rightarrow [0, 1]$ is any t-norm.
 - ▶ Connective "or" x is A or y is B use fuzzy union:
 - ▶ $A \cup B \in U \times V: \mu_{A \cup B}(x, y) = s[\mu_A(x), \mu_B(y)]$
 - ▶ $s: [0, 1] \times [0, 1] \rightarrow [0, 1]$ is any s-norm.
 - ▶ Connective "not" x is not A use fuzzy complements.
- ▶ **Example:** Recall the weather example:
 - ▶ $HO = (x$ is C and x is not H) or x is P : a fuzzy relation in the product space $[-20, 40]$:
 - ▶ $C = cold, H = hot, P = pleasant, x_1 = x_2 = x_3 = x$
 - ▶ $\mu_{HO}(x_1, x_2, x_3) = s\{t[\mu_c(x_1), c(\mu_H(x_2))], \mu_M(x_3)\}$

Interpretation of Fuzzy Rules

- ▶ For classical rules:
 - ▶ **IF p THEN q** $\rightsquigarrow p \rightarrow q$
 - ▶ $p \rightarrow q \equiv \bar{p} \vee q \equiv \bar{p} \vee (p \wedge q)$
- ▶ For fuzzy rules
 - ▶ p and q are fuzzy propositions
 - ▶ $\bar{\cdot} \equiv$ complement; $\vee \equiv$ s-norm; $\wedge \equiv$ t-norm
 - ▶ Different c-norm, s-norm, t-norm yields verity of interpretations
- ▶ **IF $\langle FP1 \rangle$ THEN $\langle FP2 \rangle$**
 - ▶ $FP1, FP2$: fuzzy relations in $U = U_1 \times \dots \times U_n, V = V_1 \times \dots \times V_m$,
 - ▶ x and y are linguistic variables (vectors) in U and V
- ▶ Some popular Fuzzy interpretation:
 - ▶ **Dienes-Rescher Implication**: Using basic fuzzy c-norm and s-norm,

 $Q_D \in U \times V$

 $\mu_{Q_D}(x, y) = \max[1 - \mu_{FP1}(x), \mu_{FP2}(y)]$

Interpretation of Fuzzy Rules

- ▶ **Lukasiewicz Implication:** Using basic fuzzy c-norm and Yager s-norm with $w = 1$, $Q_L \in U \times V$
$$\mu_{Q_L}(x, y) = \min[1, 1 - \mu_{FP1}(x) + \mu_{FP2}(y)]$$
- ▶ **Zadeh Implication:** Using basic fuzzy c-norm, s-norm, and t-norm, $Q_Z \in U \times V$
$$\mu_{Q_Z}(x, y) = \max\{\min[\mu_{FP1}(x), \mu_{FP2}(y)], 1 - \mu_{FP1}(x)\}$$
- ▶ **Gödel Implication:** a well-known implication in classical logic, $Q_G \in U \times V$
$$\mu_{Q_G}(x, y) = \begin{cases} 1 & \text{if } \mu_{FP1}(x) \leq \mu_{FP2}(y) \\ \mu_{FP2}(y) & \text{otherwise} \end{cases}$$

- ▶ **Lemma:** For all $(x, y) \in U \times V$

$$\mu_{Q_z}(x, y) \leq \mu_{Q_D}(x, y) \leq \mu_{Q_L}(x, y)$$

- ▶ **Proof:**

- ▶ $0 \leq 1 - \mu_{FP_1}(x) \leq 1$ and
 $0 \leq \mu_{FP_2}(y) \leq 1 \rightsquigarrow \max\{1 - \mu_{FP_1}(x), \mu_{FP_2}(y)\} \leq 1 - \mu_{FP_1}(x) + \mu_{FP_2}(x)$
- ▶ $\max\{1 - \mu_{FP_1}(x), \mu_{FP_2}(y)\} \leq 1 \rightsquigarrow \mu_{Q_D} \leq \mu_{Q_L}$
- ▶ $\min[\mu_{FP_1}(x), \mu_{FP_2}(y)] \leq \mu_{FP_2}(y) \rightsquigarrow \mu_{Q_z}(x, y) \leq \mu_{Q_D}(x, y)$
- ▶ We will learn later, the criteria to choose the combination of c-norms, t-norms, and s-norms for interpretation
- ▶ In crisp logic $p \rightarrow q$ is **global implication**,
 - ▶ It covers all possible choices
- ▶ In fuzzy logic $p \rightarrow q$ is **local implication**
 - ▶ Example: *IF temperature is high THEN turn on the air condition*
 - ▶ no guideline when "temperature is medium" , or "temperature is low"
- ▶ \therefore the fuzzy rule should be mentioned as:

$$IF \langle FP_1 \rangle THEN \langle FP_2 \rangle ELSE \langle NOTHING \rangle \rightsquigarrow p \rightarrow q = p \wedge q$$

- ▶ using min or algebraic product leads to **Mamdani implications** (the most popular implication)

$$\mu_{Q_{MM}}(x, y) = \min[\mu_{FP1}(x), \mu_{FP2}(y)]$$

$$\mu_{Q_{MP}}(x, y) = \mu_{FP1}(x)\mu_{FP2}(y)$$

- ▶ BUT some rule may implicitly clarify the else part
 - ▶ Example: *IF temperature is high THEN turn on the air condition*
 - ▶ It implicitly mentions that *IF temperature is low THEN turn off the air condition*
- ▶ diff. people have diff. interpretations.
- ▶ \therefore diff. implications are required
- ▶ If the human experts think that their rules are local, then use the Mamdani implications; otherwise, use the global implications

Example

- ▶ Let $U = \{-10, 5, 15, 25\}$, $V = \{1, 2, 3\}$
- ▶ $x \in U$: temperature, $y \in V$: air condition mode
- ▶ IF-THEN rule: *IF x is low THEN y is small*
- ▶ fuzzy set "low" = $1/-10 + .7/5 + .1/15 + 0/25$
- ▶ fuzzy set "small" = $1/1 + 0.5/2 + 0.1/3$
- ▶ $Q_D = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 1/(5, 1) + 0.5/(5, 2) + 0.3/(5, 3) + 1/(15, 1) + 0.9/(15, 2) + 0.9/(15, 3) + 1/(25, 1) + 1/(25, 2) + 1/(25, 3)$
- ▶ $Q_L = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 1/(5, 1) + 0.8/(5, 2) + 0.4/(5, 3) + 1/(15, 1) + 1/(15, 2) + 1/(15, 3) + 1/(25, 1) + 1/(25, 2) + 1/(25, 3)$
- ▶ $Q_Z = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 0.7/(5, 1) + 0.5/(5, 2) + 0.3/(5, 3) + 0.9/(15, 1) + 0.9/(15, 2) + 0.9/(15, 3) + 1/(25, 1) + 1/(25, 2) + 1/(25, 3)$

Example: Cont'd

- ▶ $Q_G = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 1/(5, 1) + 0.5/(5, 2) + 0.1/(5, 3) + 1/(15, 1) + 1/(15, 2) + 1/(15, 3) + 1/(25, 1) + 1/(25, 2) + 1/(25, 3)$
- ▶ $Q_{MM} = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 0.7/(5, 1) + 0.5/(5, 2) + 0.1/(5, 3) + 0.1/(15, 1) + 0.1/(15, 2) + 0.1/(15, 3) + 0/(25, 1) + 0/(25, 2) + 0/(25, 3)$
- ▶ $Q_{MM} = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 0.7/(5, 1) + 0.35/(5, 2) + 0.07/(5, 3) + 0.1/(15, 1) + 0.05/(15, 2) + 0.01/(15, 3) + 0/(25, 1) + 0/(25, 2) + 0/(25, 3)$
- ▶ membership of $(25, 1), (25, 2), (25, 3)$ is full for $Q_z, Q_D, Q_L, Q_G,$
- ▶ It is zero for Q_{MM} and Q_{MP} since $\mu_{high}(25) = 0$

- ▶ \therefore Dienes-Rescher, Lukasiewicz, Zadeh and Godel implications are global,
- ▶ Mamdani implications are local.
- ▶ **BUT** Manipulations are easier with Mamdani implications rather than others