

# Computational Intelligence

## Lecture 12: Linguistic Variables and Fuzzy Rules

Farzaneh Abdollahi

Department of Electrical Engineering


Amirkabir University of Technology

Linguistic Variables  
Linguistic Hedge

Fuzzy IF-Then Rules  
Fuzzy Proposition  
Interpretation of Fuzzy Rules

# Linguistic Variables

- ▶ **Linguistic Variable:** when a variable can take words in natural languages as its values.
  - ▶ the words are characterized by fuzzy sets defined in the universe of discourse
- ▶ **Example:** The weather temperature: a variable  $x \in [-20 \ 40]$ 
  - ▶ Let's define three fuzzy sets "Cold", "Pleasant," and "Hot"
  - ▶  $x$ : a linguistic variable,  $\rightsquigarrow$  " $x$  is cold,"
  - ▶  $x$  also can take numbers in  $[-20 \ 40] \rightsquigarrow, x = 10^\circ \text{C}$



- ▶ **Linguistic Variable:** is characterized by  $(X, T, U, M)$ 
  - ▶ **X:** the name of the linguistic variable (the weather temperature)
  - ▶ **T:** the set of linguistic values that  $X$  can take  
( $T = \{cold, pleasant, hot\}$ )
  - ▶ **U:** the actual physical domain in which the linguistic variable  $X$  takes its quantitative (crisp) values ( $U = [-20, 40]$ ).
  - ▶ **M:** a semantic rule that relates each linguistic value in  $T$  with a fuzzy set in  $U$ ; ( $M$  relates "cold," "pleasant," and "hot" with the membership functions shown in Fig)
  
- ▶ **Linguistic Hedge:**
  - ▶ One may use more than one word to describe the ling. var. ("very cold", "not pleasant", "more or less hot")
  - ▶ The value of a ling. var. is a composite of atomic term  $x = x_1 x_2 \dots x_n$
  - ▶ The atomic terms can be classified into:
    - ▶ **Primary terms:** labels of fuzzy sets; "cold," "pleasant," and "hot."
    - ▶ **Complement:** "not", "and" and "or."
    - ▶ **Hedges:** "very," "slightly," "more or less,"

## Some Examples of Linguistic Hedge

- ▶ Let  $A$  be a fuzzy set in  $U$ 
  - ▶ **very  $A$**  : a fuzzy set in  $U$   $\mu_{\text{very } A}(x) = [\mu_A(x)]^2$
  - ▶ **more or less  $A$** : a fuzzy set in  $U$   $\mu_{\text{more or less } A}(x) = [\mu_A(x)]^{1/2}$
- ▶ **Example:**  $U = \{10, \dots, 130\}$ 
  - ▶ fuzzy set "fast" =  $1/130 + 1/120 + 0.8/100 + 0.6/80 + 0.1/40$
  - ▶ "very very fast" =  $1/130 + 1/120 + 0.4096/100 + 0.1296/80 + 0.0001/40$
  - ▶ "more or less fast" =  $1/130 + 1/120 + 0.8944/100 + 0.7746/80 + 0.3162/40$

# Fuzzy IF-Then Rules

- ▶ **IF** <fuzzy proposition>, **THEN** <fuzzy proposition>
- ▶ There are two types of fuzzy proposition
  - ▶ **Atomic fuzzy proposition**: a single statement  
 $x$  is  $A$
  - ▶ **Compound fuzzy proposition**: a composition of atomic fuzzy propositions using the connectives "and," "or," and "not"  
 $x$  is not  $S$  and  $x$  is not  $F$
  - ▶  $S, F, A$  are fuzzy sets
- ▶ Compound fuzzy propositions should be considered as fuzzy relations.

- ▶ Let  $x, y$ : linguistic variables in the physical domains  $U, V$ , and  $A, B$ : fuzzy sets in  $U, V$ 
  - ▶ Connective "and"  $x$  is  $A$  and  $y$  is  $B$  use fuzzy intersection:
    - ▶  $A \cap B \in U \times V: \mu_{A \cap B}(x, y) = t[\mu_A(x), \mu_B(y)]$
    - ▶  $t: [0, 1] \times [0, 1] \rightarrow [0, 1]$  is any t-norm.
  - ▶ Connective "or"  $x$  is  $A$  or  $y$  is  $B$  use fuzzy union:
    - ▶  $A \cup B \in U \times V: \mu_{A \cup B}(x, y) = s[\mu_A(x), \mu_B(y)]$
    - ▶  $s: [0, 1] \times [0, 1] \rightarrow [0, 1]$  is any s-norm.
  - ▶ Connective "not"  $x$  is not  $A$  use fuzzy complements.

- ▶ Let  $x, y$ : linguistic variables in the physical domains  $U, V$ , and  $A, B$ : fuzzy sets in  $U, V$ 
  - ▶ Connective "and"  $x$  is  $A$  and  $y$  is  $B$  use fuzzy intersection:
    - ▶  $A \cap B \in U \times V$ :  $\mu_{A \cap B}(x, y) = t[\mu_A(x), \mu_B(y)]$
    - ▶  $t : [0, 1] \times [0, 1] \rightarrow [0, 1]$  is any t-norm.
  - ▶ Connective "or"  $x$  is  $A$  or  $y$  is  $B$  use fuzzy union:
    - ▶  $A \cup B \in U \times V$ :  $\mu_{A \cup B}(x, y) = s[\mu_A(x), \mu_B(y)]$
    - ▶  $s : [0, 1] \times [0, 1] \rightarrow [0, 1]$  is any s-norm.
  - ▶ Connective "not"  $x$  is not  $A$  use fuzzy complements.
- ▶ **Example:** Recall the weather example:
  - ▶  $HO = (x$  is  $C$  and  $x$  is not  $H$ ) or  $x$  is  $P$ : a fuzzy relation in the product space  $[-20, 40]$ :
  - ▶  $C = \text{cold}, H = \text{hot}, P = \text{pleasant}, x_1 = x_2 = x_3 = x$
  - ▶  $\mu_{HO}(x_1, x_2, x_3) = s\{t[\mu_c(x_1), c(\mu_H(x_2))], \mu_M(x_3)\}$



# Interpretation of Fuzzy Rules

- ▶ For classical rules:
  - ▶ **IF  $p$  THEN  $q$**   $\rightsquigarrow p \rightarrow q$
  - ▶  $p \rightarrow q \equiv \bar{p} \vee q \equiv \bar{p} \vee (p \wedge q)$
- ▶ For fuzzy rules
  - ▶  $p$  and  $q$  are fuzzy propositions
  - ▶  $\bar{\cdot} \equiv$  complement;  $\vee \equiv$  s-norm;  $\wedge \equiv$  t-norm
  - ▶ Different c-norm, s-norm, t-norm yields verity of interpretations
- ▶ **IF  $\langle FP1 \rangle$  THEN  $\langle FP2 \rangle$** 
  - ▶  $FP1, FP2$ : fuzzy relations in  $U = U_1 \times \dots \times U_n, V = V_1 \times \dots \times V_m$ ,
  - ▶  $x$  and  $y$  are linguistic variables (vectors) in  $U$  and  $V$
- ▶ Some popular Fuzzy interpretation:
  - ▶ **Dienes-Rescher Implication**: Using basic fuzzy c-norm and s-norm,
   
 $Q_D \in U \times V$ 
  
 $\mu_{Q_D}(x, y) = \max[1 - \mu_{FP1}(x), \mu_{FP2}(y)]$

# Interpretation of Fuzzy Rules

- ▶ **Lukasiewicz Implication:** Using basic fuzzy c-norm and Yager s-norm with  $w = 1$ ,  $Q_L \in U \times V$   
$$\mu_{Q_L}(x, y) = \min[1, 1 - \mu_{FP1}(x) + \mu_{FP2}(y)]$$
- ▶ **Zadeh Implication:** Using basic fuzzy c-norm, s-norm, and t-norm,  $Q_Z \in U \times V$   
$$\mu_{Q_Z}(x, y) = \max\{\min[\mu_{FP1}(x), \mu_{FP2}(y)], 1 - \mu_{FP1}(x)\}$$
- ▶ **Gödel Implication:** a well-known implication in classical logic,  $Q_G \in U \times V$   
$$\mu_{Q_G}(x, y) = \begin{cases} 1 & \text{if } \mu_{FP1}(x) \leq \mu_{FP2}(y) \\ \mu_{FP2}(y) & \text{otherwise} \end{cases}$$

- ▶ **Lemma:** For all  $(x, y) \in U \times V$   

$$\mu_{Q_z}(x, y) \leq \mu_{Q_D}(x, y) \leq \mu_{Q_L}(x, y)$$

- ▶ **Proof:**

- ▶  $0 \leq 1 - \mu_{FP_1}(x) \leq 1$  and  
 $0 \leq \mu_{FP_2}(y) \leq 1 \rightsquigarrow \max\{1 - \mu_{FP_1}(x), \mu_{FP_2}(y)\} \leq 1 - \mu_{FP_1}(x) + \mu_{FP_2}(x)$
- ▶  $\max\{1 - \mu_{FP_1}(x), \mu_{FP_2}(y)\} \leq 1 \rightsquigarrow \mu_{Q_D} \leq \mu_{Q_L}$
- ▶  $\min[\mu_{FP_1}(x), \mu_{FP_2}(y)] \leq \mu_{FP_2}(y) \rightsquigarrow \mu_{Q_z}(x, y) \leq \mu_{Q_D}(x, y)$
- ▶ We will learn later, the criteria to choose the combination of c-norms, t-norms, and s-norms for interpretation
- ▶ In crisp logic  $p \rightarrow q$  is **global implication**,
  - ▶ It covers all possible choices
- ▶ In fuzzy logic  $p \rightarrow q$  is **local implication**
  - ▶ Example: *IF temperature is high THEN turn on the air condition*
  - ▶ no guideline when "temperature is medium" , or "temperature is low"
- ▶  $\therefore$  the fuzzy rule should be mentioned as:  

$$IF \langle FP_1 \rangle THEN \langle FP_2 \rangle ELSE \langle NOTHING \rangle \rightsquigarrow p \rightarrow q = p \wedge q$$

- ▶ using min or algebraic product leads to **Mamdani implications** (the most popular implication)

$$\mu_{Q_{MM}}(x, y) = \min[\mu_{FP1}(x), \mu_{FP2}(y)]$$

$$\mu_{Q_{MP}}(x, y) = \mu_{FP1}(x)\mu_{FP2}(y)$$

- ▶ BUT some rule may implicitly clarify the else part
  - ▶ Example: *IF temperature is high THEN turn on the air condition*
  - ▶ It implicitly mentions that *IF temperature is low THEN turn off the air condition*
- ▶ diff. people have diff. interpretations.
- ▶  $\therefore$  diff. implications are required
- ▶ If the human experts think that their rules are local, then use the Mamdani implications; otherwise, use the global implications

## Example

- ▶ Let  $U = \{-10, 5, 15, 25\}$ ,  $V = \{1, 2, 3\}$
- ▶  $x \in U$ : temperature,  $y \in V$ : air condition mode
- ▶ IF-THEN rule: *IF  $x$  is low THEN  $y$  is small*
- ▶ fuzzy set "low" =  $1/-10 + .7/5 + .1/15 + 0/25$
- ▶ fuzzy set "small" =  $1/1 + 0.5/2 + 0.1/3$
- ▶  $Q_D = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 1/(5, 1) + 0.5/(5, 2) + 0.3/(5, 3) + 1/(15, 1) + 0.9/(15, 2) + 0.9/(15, 3) + 1/(25, 1) + 1/(25, 2) + 1/(25, 3)$
- ▶  $Q_L = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 1/(5, 1) + 0.8/(5, 2) + 0.4/(5, 3) + 1/(15, 1) + 1/(15, 2) + 1/(15, 3) + 1/(25, 1) + 1/(25, 2) + 1/(25, 3)$
- ▶  $Q_Z = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 0.7/(5, 1) + 0.5/(5, 2) + 0.3/(5, 3) + 0.9/(15, 1) + 0.9/(15, 2) + 0.9/(15, 3) + 1/(25, 1) + 1/(25, 2) + 1/(25, 3)$

## Example: Cont'd

- ▶  $Q_G = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 1/(5, 1) + 0.5/(5, 2) + 0.1/(5, 3) + 1/(15, 1) + 1/(15, 2) + 1/(15, 3) + 1/(25, 1) + 1/(25, 2) + 1/(25, 3)$
- ▶  $Q_{MM} = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 0.7/(5, 1) + 0.5/(5, 2) + 0.1/(5, 3) + 0.1/(15, 1) + 0.1/(15, 2) + 0.1/(15, 3) + 0/(25, 1) + 0/(25, 2) + 0/(25, 3)$
- ▶  $Q_{MM} = 1/(-10, 1) + 0.5/(-10, 2) + 0.1/(-10, 3) + 0.7/(5, 1) + 0.35/(5, 2) + 0.07/(5, 3) + 0.1/(15, 1) + 0.05/(15, 2) + 0.01/(15, 3) + 0/(25, 1) + 0/(25, 2) + 0/(25, 3)$
- ▶ membership of  $(25, 1), (25, 2), (25, 3)$  is full for  $Q_z, Q_D, Q_L, Q_G,$
- ▶ It is zero for  $Q_{MM}$  and  $Q_{MP}$  since  $\mu_{high}(25) = 0$

- ▶  $\therefore$  Dienes-Rescher, Lukasiewicz, Zadeh and Godel implications are global,
- ▶ Mamdani implications are local.
- ▶ **BUT** Manipulations are easier with Mamdani implications rather than others