

Computational Intelligence

Lecture 3:Fuzzy Relations

Farzaneh Abdollahi

Department of Electrical Engineering

Amirkabir University of Technology

Fall 2010

Classical Relation

Fuzzy Relation

- Projection

- Cylindrical Extension

- Cartesian Product of Fuzzy Sets

- Composition

Extension Principle

3/22

Example:

- ▶ $U = \{1, 2, 3\}$, $V = \{a, b\}$
- ▶ $U \times V = (1, a), (1, b), (2, a), (2, b), (3, a), (3, b)$
- ▶ let $Q(U, V)$ be a relation named "the first element is no smaller than 2"

- ▶ $Q(U, V) = \{(2, a), (2, b), (3, a), (3, b)\}$


$U \setminus V$	a	b
1	0	0
2	1	1
3	1	1

Example: Dormitory based on Distance of cities.

- ▶ $V = \{Tehran, Tabriz, Karaj, Qom\}$, $U = \{Tehran, Esfahan\}$
- ▶ Relation: "very far"
- ▶ use number between 0 and 1 for degree of relation

U/V	Tehran	Tabriz	Karaj	Qom
Tehran	0	0.9	0.1	0.3
Esfahan	0.7	0.95	0.8	0.5


Projection



Projection

► **Example:** A crisp relation in $V \times U = R^2$

- $A = \{(x, y) \in R^2 \mid (x-1)^2 + (y-1)^2 \leq 1\}$
- A_1 the projection of A on U : $[0, 1] \subset U$
- A_2 the projection of A on V : $[0, 1] \subset V$



Projection of Fuzzy Sets

- ▶ Q : a fuzzy relation in $U_1 \times \dots \times U_n$
- ▶ $\{i_1, \dots, i_k\}$ a subsequence of $\{1, 2, \dots, n\}$
- ▶ The projection of Q on $U_{i_1} \times \dots \times U_{i_k}$ is a fuzzy relation Q_P in $U_{i_1} \times \dots \times U_{i_k}$ s.t.:

$$\mu_{Q_P}(u_{i_1}, \dots, u_{i_k}) = \max_{u_{j_1} \in U_{j_1}, \dots, u_{j_{n-k}} \in U_{j_{n-k}}} \mu_Q(u_1, \dots, u_n)$$

- ▶ $\{u_{j_1}, \dots, u_{j_{n-k}}\}$ is the complement of $\{u_{i_1}, \dots, u_{i_k}\}$
- ▶ For binary fuzzy relation $U \times V$:

- ▶ Q_1 projection in U : $\mu_{Q_1}(x) = \max_{y \in V} \mu_Q(x, y)$

- ▶ **Example:** Recall the "AE" example

- ▶ Projection in U : $AE_1 = \int_U \max_{y \in V} e^{-(x-y)^2} / x = \int_U 1/x$
- ▶ Projection in V : $AE_2 = \int_V \max_{x \in U} e^{-(x-y)^2} / y = \int_V 1/y$


- ▶ **Example:** Recall the "very far" example

- ▶ Q_1 : projection on u : $Q_1 = 0.9 / \text{Tehran} + 0.95 / \text{Esfahan}$
- ▶ Q_2 : projection on V :

$$Q_2 = 0.7 / \text{Tehran} + 0.95 / \text{Tabriz} + 0.8 / \text{Karaj} + 0.5 / \text{Qom}$$

Cylindrical Extension

- ▶ Extending the projection of fuzzy cylindrically.
- ▶ **Example:** Recall the circle example
 - ▶ A_{1E} Cylindric extension of A_1 to $U \times V = R^2$
 - ▶ $A_{1E} = [0, 1] \times (-\infty, \infty) \subset R^2$
- ▶ The projection constrains a fuzzy relation to a subspace
- ▶ The cylindric extension extends a fuzzy relation (or fuzzy set) from a subspace to the whole space.



- Let Q_p be a fuzzy relation in $U_{i_1} \times \dots \times U_{i_k}$ and $\{i_1, \dots, i_k\}$ is a subsequence of $\{1, 2, \dots, n\}$, then the **cylindric extension** of Q_p to $U_1 \times \dots \times U_n$ is a fuzzy relation Q_{pE} in $U_1 \times \dots \times U_n$

$$\mu_{Q_{pE}}(u_1, \dots, u_n) = \mu_{Q_p}(u_{i_1}, \dots, u_{i_k})$$

- ▶ For binary set:
 - ▶ $U \times V$,
 - ▶ Q_1 a fuzzy set in U
 - ▶ Q_{1E} the cylindric extension to $U \times V$
 - ▶ $\mu_{Q_{1E}}(x, y) = \mu_{Q_1}(x)$
- ▶ **Example:** Recall the "AE" example
 - ▶ $A_{E_{1E}} = \int_{U \times V} 1/(x, y) = U \times V$
 - ▶ $A_{E_{2E}} = \int_{U \times V} 1/(x, y) = U \times V$


Cartesian Product of Fuzzy Sets

- Let $A_1, \dots, A_n$ be fuzzy sets in $U_1, \dots, U_n$, respectively. The Cartesian product of $A_1, \dots, A_n$ denoted by $A_1 \times \dots \times A_n$, is a fuzzy relation in $U_1 \times \dots \times U_n$:

$$\mu_{A_1 \times \dots \times A_n}(u_1, \dots, u_n) = \mu_{A_1}(u_1) * \dots * \mu_{A_n}(u_n)$$

- where $*$ represents any t-norm operator.
- Lemma:** If Q is a fuzzy relation in $U_1 \times \dots \times U_n$ and $Q_1, \dots, Q_n$ are its projections on $U_1, \dots, U_n$, respectively, then

$$Q \subset Q_1 \times \dots \times Q_n$$




where we use "min" for the t-norm

Composition

- ▶ $P(U, V)$ and $Q(V, W)$: two crisp binary relations that share a common set V .
- ▶ The composition of P and Q , $(P \circ Q)$, is a relation in $U \times W$ s.t. $(x, z) \in Q$ iff there exists at least one $y \in V$ s.t. $(x, y) \in P$ and $(y, z) \in Q$.
- ▶ **Lemma:** $P \circ Q$ is the composition of $P(U, V)$ and $Q(V, W)$ iff

$$\mu_{P \circ Q}(x, z) = \max_t [\mu_P(x, y), \mu_Q(y, z)] \quad (1)$$

for any $(x, z) \in U \times W$, where t is any t -norm.



► Proof:

- ▶ If PoQ is the composition:
 - ▶ $(x, z) \in PoQ \rightsquigarrow \exists y \in V$ s.t. $\mu_P(x, y) = 1 \& \mu_Q(y, z) = 1$
 - ▶ $\therefore \mu_{PoQ}(x, z) = 1 = \max_{y \in V} t[\mu_P(x, y), \mu_Q(y, z)]$
 - ▶ If $(x, z) \notin PoQ \rightsquigarrow$ for any $y \in V$, $\mu_P(x, y) = 0$ or $\mu_Q(y, z) = 0$
 - ▶ $\therefore \mu_{PoQ}(x, z) = 0 = \max_{y \in V} t[\mu_P(x, y), \mu_Q(y, z)]$.
 - ▶ Eq. (1) is true.
- ▶ Conversely, if the Eq. (1) is true:
 - ▶ $(x, z) \in PoQ \rightsquigarrow \max_{y \in V} t[\mu_P(x, y), \mu_Q(y, z)] = 1$
 - ▶ $\therefore$ there exists at least one $y \in V$ s.t. $\mu_P(x, y) = \mu_Q(y, z) = 1$ (Axiom t1)
 - ▶ For $(x, z) \notin PoQ \rightsquigarrow \max_{y \in V} [\mu_P(x, y), \mu_Q(y, z)] = 0$
 - ▶ $\therefore \nexists y \in V$ s.t. $\mu_P(x, y) = \mu_Q(y, z) = 1$.
 - ▶ $\therefore PoQ$ is the composition

Fuzzy Composition

- ▶ Composition for fuzzy relations is defined similar to crisp relations
- ▶ Based on different definition of t-norm different composition is obtained.
- ▶ The two most popular compositions:
 - ▶ **Max-Min**: of fuzzy relations $P(U, V)$ and $Q(V, W)$ is a fuzzy relation $P \circ Q$ in $U \times W$ s.t. $\mu_{P \circ Q}(x, z) = \max_{y \in V} \min[\mu_P(x, y), \mu_Q(y, z)]$
 - ▶ It uses the min for t-normwhere $(x, z) \in U \times W$.
 - ▶ **Max-Product**: of fuzzy relations $P(U, V)$ and $Q(V, W)$ is a fuzzy relation $P \circ Q$ in $U \times W$ s.t. $\mu_{P \circ Q}(x, z) = \max_{y \in V} [\mu_P(x, y) \cdot \mu_Q(y, z)]$ where $(x, z) \in U \times W$.
 - ▶ It uses algebraic product for t-norm

Example: Recall Dormitory example

- $V = \{Tehran, Tabriz, Karaj, Qom\}$, $U = \{Tehran, Esfahan\}$, $W = \{Boomehen, Kashan, Ardebil\}$

- $P(U, V)$ "very far"

U/V	Tehran	Tabriz	Karaj	Qom
Tehran	0	0.9	0.1	0.3
Esfahan	0.7	0.95	0.8	0.5

Example: Recall Dormitory example

- $V = \{Tehran, Tabriz, Karaj, Qom\}$, $U = \{Tehran, Esfahan\}$, $W = \{Boomehen, Kashan, Ardebil\}$

- $P(U, V)$ "very far"

U/V	Tehran	Tabriz	Karaj	Qom
Tehran	0	0.9	0.1	0.3
Esfahan	0.7	0.95	0.8	0.5

- $Q(V, W)$:"very near"

V/W	Boomehen	Kashan	Ardebil
Tehran	1	0.4	0.1
Tabriz	0.2	0	0.8
Karaj	0.6	0.3	0.1
Qom	0.7	0.95	0

► $PoQ(U, W)$ using Max-min

P

U/V	Tehran	Tabriz	Karaj	Qom
Tehran	0	0.9	0.1	0.3
Esfahan	0.7	0.95	0.8	0.5

Q

$V \setminus W$	Boomehen	Kashan	Ardebil
Tehran	1	0.4	0.1
Tabriz	0.2	0	0.8
Karaj	0.6	0.3	0.1
Qom	0.7	0.95	0

Min

0	0.9	0.1	0.3
1	0.2	0.6	0.7
0	0.2	0.1	0.3

Max

PoQ

U/W	Boomehen	Kashan	Ardebil
Tehran	0.3	0.3	0.8
Esfahan	0.7	0.5	0.8

► $PoQ(U, W)$ using Max-Product

P

U/V	Tehran	Tabriz	Karaj	Qom
Tehran	0	0.9	0.1	0.3
Esfahan	0.7	0.95	0.8	0.5

Q

$V \setminus W$	Boomehen	Kashan	Ardebil
Tehran	1	0.4	0.1
Tabriz	0.2	0	0.8
Karaj	0.6	0.3	0.1
Qom	0.7	0.95	0

x

0	0.9	0.1	0.3
1	0.2	0.6	0.7
0	0.18	0.06	0.21

Max

PoQ

U/W	Boomehen	Kashan	Ardebil
Tehran	0.21	0.0285	0.72
Esfahan	0.7	0.475	0.76

- ▶ The relational matrix for the fuzzy composition $P \circ Q$ can be computed according to the following method:
 - ▶ For max-min composition
 - ▶ write out each element in the matrix product PQ , But treat:
 - ▶ each multiplication as a min operation
 - ▶ each addition as a max operation
 - ▶ For max-product composition,
 - ▶ write out each element in the matrix product PQ , but treat
 - ▶ each addition as a max operation.

Extension Principle

- ▶ **Extension Principle:** $\mu_B(y) = \max_{x \in f^{-1}(y)} \mu_A(x), y \in V$
 - ▶ $f^{-1}(y)$: set of all points $x \in U$ s.t. $f(x) = y$
- ▶ **Example** $U = \{1, \dots, 10\}, x \in U, f(x) = x^2 \in V = \{1, \dots, 100\}$
- ▶ Fuzzy set: "small" = $1/1 + 1/2 + 0.8/3 + 0.6/4 + 0.4/5$
- ▶ $\therefore$ "small"² = $1/1 + 1/4 + 0.8/9 + 0.6/16 + 0.4/25$